

## PAPILIO PHORCAS.

\*\*\*\*\*

### CHARACTER GENERICUS.

*Antennæ apicem versus crassiores, sæpius clavato-capitatae.*

*Alæ (sedentis) erectæ sursumque conniventes,  
(volatu diurno.)*

*Lin. Syst. Nat. p. 744.*

### CHARACTER SPECIFICUS, &c.

PAPILIO niger, fascia supra viridi subtus pallidiore, alis posterioribus caudatis.

PAPILIO Doreus, P. alis fascia supra viridi, subtus alba.

*Lin. Syst. Nat. Gmel. p. 2239.*

*Fab. sp. ins. 2. p. 18.*

## PAPILIO PHORCAS.

*Cram. 1. t. 2. f. b. c.*

---

In Sierra Leona præcipue conspicitur Papilio Phorcus, inter pulchriores merito numerandus. Magnitudinem naturalem ostendit tabula.





J



456

## PHORCAS.

---

### *GENERIC CHARACTER.*

*Antennæ* gradually thickening towards the end,  
terminating in a clavated tip.

*Wings*, when at rest, upright. Flight diurnal.

### *SPECIFIC CHARACTER.*

Black Butterfly, with green band, paler beneath,  
the lower wings tailed.

---

This species, which may justly be numbered among  
the most elegant of its tribe, is chiefly found in Sierra  
Leona, and is represented on the plate in its natural  
size.